

Stars older than the universe and possible mechanism of their creation

A. D. Dolgov^{a,b,c*}

^a *University of Ferrara
Ferrara 40100, Italy*

^b *Novosibirsk State University
Novosibirsk, 630090, Russia*

^c *ITEP
Moscow, 117218, Russia*

The only aspect of LCDMHC cosmology that is agreeable to HGD cosmology is that the universe began with a big bang event. CHG

Abstract

An impressive bulk of multiple astronomical observations indicates that there are plenty of objects in the universe with the age which cannot be explained by the conventional theory. A model is considered which successfully describes all these puzzling phenomena.

1 Introduction

Astronomical observations demonstrate the general trend, indicating that many objects in the universe were formed much earlier than expected by theory and possibly so old objects are even not allowed by the standard theory. Among them there are stars in the Milky Way, older than the Galaxy and even older than the universe, within at least two sigma, distant high redshift ($z \sim 10$) objects. such as early galaxies, QSO/supermassive BHs, gamma-bursters, and early supenovae.

From HGD cosmology, the first stars form at $\sim 300,000$ yrs + 30,000 yrs.

If no explanation is found in the conventional frameworks, there are two possible ways (maybe more) to explain these puzzling phenomena:

1. A novel mechanism of formation of stellar-type objects in very early universe [1].
2. Modification of the cosmological expansion regime in such a way that the universe becomes older than calculated in the frameworks of the standard model, as is done e.g. in ref. [2].

Here we discuss the first possibility, more detail of which can be found in our paper [1]. First, let us present the expression of the universe age t_U , as a function of the cosmological redshift z :

$$t(z) = \frac{1}{H} \int_0^{1/(z+1)} \frac{dx}{\sqrt{1 - \Omega_{tot} + \Omega_m/x + x^2 \Omega_v}}, \quad (1)$$

where Ω_a is the fractional energy density of matter (Ω_m , baryonic plus dark matter), of dark energy (Ω_v), and of the total cosmological energy density (Ω_{tot}). According to the Planck data, the present day values of these parameters are: $\Omega_{tot} = 1$, $\Omega_m = 0.317$, and $\Omega_v = 0.683$. There is some tension between the values of the Hubble parameter measured by Planck, $H_{pl} = 67.3$ km/sec/Mpc and by the traditional astronomical methods, which can lead to H as large as

* e-mail: dolgov@fe.infn.it

Kinematic viscosity of the plasma epoch is too high for objects smaller than galaxies to form.

$H_{astr} = 74$ km/sec/Mpc, see ref. [3] for discussion. We present a few examples of the universe age in gigayears for different z , the first number corresponds to the Planck value of H , and the other, shorter one, in brackets to the larger astronomical value: $t_U \equiv t(0) = 13.8$ (12.5.); $t(3) = 2.14$ (1.94); $t(6) = 0.93$; (0.82); $t(10) = 0.47$ (0.43); and $t(12) = 0.37$; (0.33).

2 Old stars in the Milky Way

True, so LCDMHC fails, not the big bang.

Recently several stars have been discovered in the Galaxy which ages are unexpectedly high.

Employing thorium and uranium abundances in comparison with each other and with several stable elements the age of metal-poor, halo star BD+17° 3248 was estimated as 13.8 ± 4 Gyr, as is argued in ref. [4]. For comparison the estimated age of the inner halo of the Galaxy is 11.4 ± 0.7 Gyr [5]

The age of a star in the galactic halo, HE 1523-0901, was estimated to be about 13.2 Gyr [6]. In this work many different chronometers, such as the U/Th, U/Ir, Th/Eu, and Th/Os ratios to measure the star age have been employed for the first time.

Most puzzling is probably the determination of the age of metal deficient high velocity sub-giant in the solar neighborhood, HD 140283, which has the age 14.46 ± 0.31 Gyr [7]. The central value of the age exceeds the universe age by two standard deviations, if $H = 67.3$ km/sec/Mpc, and for larger $H = 74$ km/sec/Mpc the star is older than the universe more than by six standard deviations.

HGD cosmology does not constrain age of stars by velocity.

3 High redshift distant objects

3.1 Galaxies

Galaxies at high redshifts, $z \sim 10$, cannot be observed with the usual optical telescopes, which are not sensitive enough for such distant objects. Fortunately natural gravitational lens telescopes allow to see them, if the "telescope" happens to be on the light ray from the galaxy to terrestrial observers. In such a way a galaxy at $z \approx 9.6$ was discovered [8]. The galaxy was formed when the universe was about 500 million years old.

Even more striking, a galaxy at $z \approx 11$ has been observed [9] which was formed before the universe age was 0.41 Gyr (or even shorter with larger H).

Star formation is inhibited in interstellar gas with low metallicity, because higher fraction of metals enhances gas cooling and facilitates gravitational capture. Interstellar medium is enriched by metals through supernova explosion. So we need either abundant very early supervonae, or an unusual source of metallicity, and a new mechanism of galaxy formation. To make things more puzzling some observed early galaxies are indeed enriched with metals, see subsection 3.3.

Quoting ref. [10]: "Observations with WFC3/IR on the Hubble Space Telescope and the use of gravitational lensing techniques have facilitated the discovery of galaxies as far back as z 10-12, a truly remarkable achievement. However, this rapid emergence of high- z galaxies, barely 200 Myr after the transition from Population III star formation to Population II, appears to be in conflict with the standard view of how the early Universe evolved." - the quotation to be continued on top of the next subsection.

3.2 Quasars and supermassive black holes at high z and now

Continuing the quotation from ref. [10]: "This problem is very reminiscent of the better known (and probably related) premature appearance of supermassive black holes at $z \sim 6$. It is difficult

to understand how $10^9 M_\odot$ black holes appeared so quickly after the big bang without invoking non-standard accretion physics and the formation of massive seeds, both of which are not seen in the local Universe.”

A quasar with maximum $z = 7.085$ has been discovered [11], i.e. it was formed at $t < 0.75$ Gyr. Its luminosity is $6.3 \cdot 10^{13} L_\odot$ and mass $2 \cdot 10^9 M_\odot$. The quasars are supposed to be supermassive black holes (BH) and their formation in such short time looks problematic by conventional mechanisms.

There are strong indications that every large galaxy, as well as some relatively small ones, contain central supermassive black hole. The mass of the black hole may be larger than ten billions M_\odot in giant elliptical and compact lenticular galaxies and about a few million M_\odot in spiral galaxies like Milky Way. The mass of BH is typically 0.1% of the mass of the stellar bulge of galaxy, but some galaxies may have huge BH: e.g. NGC 1277 has the central BH of $1.7 \times 10^{10} M_\odot$, or 60% of its bulge mass [12]. Another interesting example is a possible existence of a supermassive black hole in an Ultracompact Dwarf Galaxy, M60-UCD1 [13] with the mass of about 20 million solar mass, which is 15% of the object’s total mass. According to the conclusion of the authors, the high black hole mass and mass fraction suggest that M60-UCD1 is the stripped nucleus of a galaxy. On the other hand, the authors observed ”that M60-UCD1’s stellar mass is consistent with its luminosity, implying many other UCDs may also host supermassive black holes. This suggests a substantial population of previously unnoticed supermassive black holes.”

These facts create serious problems for the standard scenario of formation of central supermassive BHs by accretion of matter in the central part of a galaxy. An inverted picture looks more plausible, when first a supermassive black holes were formed and attracted matter serving as seeds for subsequent galaxy formation.

3.3 Early Supernovae

The medium around the observed early quasars contains considerable amount of “metals” (elements heavier than He). According to the standard picture, only elements up to ^4He and traces of Li, Be, B were formed in the early universe by BBN, while heavier elements were created by stellar nucleosynthesis and dispersed in the interstellar space by supernova explosions. If so, prior to QSO creation a rapid star formation should take place. These stars had to produce plenty of supernovae which might enrich interstellar space by metals.

Observations of high redshift gamma ray bursters (GBR) also indicate a high abundance of supernova at large redshifts, if GBRs are very early supernovae. The highest redshift of the observed GBR is 9.4 [14] and there are a few more GBRs with smaller but still high redshifts. The necessary star formation rate for explanation of these early GBRs is at odds with the canonical star formation theory.

A recent discovery [15] of an ultra-compact dwarf galaxy older than 10 Gyr, enriched with metals, and probably with a massive black hole in its center seems to be at odds with the standard model as well. The dynamical mass of this galaxy is $2 \times 10^8 M_\odot$ and its radius is $R \sim 24$ pc, so the galaxy density is extremely high. There is a variable central X-ray source with luminosity $L_X \sim 10^{38}$ erg/s, which may be an AGN associated with a massive black hole or a low-mass X-ray binary.

4 A model of formation of compact stellar-like objects and heavy PBH in the very early universe

Quite probably the described above puzzling existence of very old objects in the early high metallicity universe will find an explanation in the frameworks of the conventional astrophysics. However, in absence of such explanation a search for mechanisms based on new physics is also desirable. We present here an explanation based on early works [16], where a simple generalization of the well known Affleck-Dine [17] scenario of baryogenesis allows to explain all observational data described above.

The modification of the Affleck-Dine (AD) scenario of baryogenesis, which give rise to significant production of stellar-like objects or heavy primordial black holes can be achieved by a simple addition of a general renormalizable coupling of the scalar baryon, χ , to the inflaton field, Φ :

$$U(\chi, \Phi) = U_\chi(\chi) + U_\Phi(\Phi) + U_{\text{int}}(\chi, \Phi). \quad (2)$$

Here $U_\Phi(\Phi)$ is the inflaton potential, $U_\chi(\chi)$ is the quartic Affleck-Dine potential, which generically has some flat directions (valleys). The potential has the form:

$$U_\chi(\chi) = [m_\chi^2 \chi^2 + \lambda_\chi (\chi^4 + |\chi|^4) + h.c.] + \lambda_2 |\chi|^4 \ln \frac{|\chi|^2}{\sigma^2}, \quad (3)$$

where the last term is the Coleman-Weinberg correction [18], which arises as a result of summation of one-loop diagrams in scalar field theory with quartic interaction term.

In the classical AD-scenario field χ acquires a large expectation value along a flat direction, e.g. during inflation and evolves down later, when the Hubble parameter dropped below m_χ . If flat directions in quadratic and quartic parts of the potential do not coincide, then at the approach to the minimum χ starts to "rotate" in two dimensional $\{Re\chi, Im\chi\}$ -plane. Rotation means that χ acquires (a large) average baryonic number.

The additional interaction term of χ with the inflaton, Φ , is taken in the form:

$$U_{\text{int}}(\chi, \Phi) = \lambda_1 |\chi|^2 (\Phi - \Phi_1)^2, \quad (4)$$

where Φ_1 is some value of the inflaton field which it passes during inflation and λ_1 is a constant. So there is a mild tuning, but otherwise this is general renormalizable coupling between χ and Φ , which surely must exist. This terms acts as a positive time-dependent mass and thus it almost always kept the gate to the valleys closed, except for a short period when when Φ is near Φ_1 . So there is a small chance for χ to reach a high value and to create a large baryon asymmetry. The behavior of the potential $U_\chi(\chi) + U_{\text{int}}(\chi, \Phi)$ for different values of the effective mass is presented in fig. 1. The potential evolves down from the upper to the lower curve reaching the latter when $\Phi = \Phi_1$ and then the potential returns back to the higher curve, when Φ drops below Φ_1 .

Correspondingly field χ rolls down toward the deeper minimum, oscillates there following the evolution of the minimum, rolls back to the origin, and starts to rotate around it, as is shown in fig. 2.

Since the inflaton opens the gate to the deeper minimum only for a short time, the probability for χ to reach the high value is low, so in most of space baryogenesis creates normal tiny baryon asymmetry, but in some bubbles which occupy a small fraction of the whole volume, baryon asymmetry may be huge.

After the QCD phase transition, the contrast in baryonic charge density transformed into perturbations of the energy/mass density and the bubbles with high B formed PBH's or compact stellar-like objects. The mass distribution of these high-B bubbles has practically model




Figure 1: Behavior of $U_\chi(\chi)$ for different values of $m_{eff}^2(t)$.




Figure 2: Evolution of $|\chi|$ with time.

independent form:

$$\frac{dN}{dM} = C_M \exp \left[-\gamma \ln^2 \frac{(M - M_1)^2}{M_0^2} \right]. \quad (5)$$

with the model dependent parameters C_M , M_1 , and M_0 .

The values of the parameters can be adjusted in such a way that superheavy BHs formed at the tail of this distribution would be abundant enough to be present in every large galaxy and in some small ones. Such heavy PBHs could be seeds for the galaxy formation. As we mentioned above, there is no satisfactory mechanism of creation of superheavy black holes in the frameworks of the usual physics, while the considered here mechanism can successfully achieve that.

This mass distribution naturally explains some unusual features of stellar mass black holes in the Galaxy. It was found that their masses are concentrated in narrow range $(7.8 \pm 1.2)M_\odot$ [19]. This result agrees with another paper where a peak around $8M_\odot$, a paucity of sources with masses below $5M_\odot$, and a sharp drop-off above $10M_\odot$ are observed [20]. These features are not explained in the standard model.

A modifications of U_{int} leads to a more complicated mass spectrum of the early formed stellar type objects, e.g., if:

$$U_{int} = \lambda_1 |\chi|^2 (\Phi - \Phi_1)^2 (\Phi - \Phi_2)^2, \quad (6)$$

we come to a two-peak mass distribution of the PBHs and compact stars, which is probably observed [21], but not yet explained.

Evolved chemistry in the so early formed QSOs can be explained, at least to some extent, by more efficient production of metals during BBN due to much larger ratio $\beta = N_B/N_\gamma$. The standard BBN essentially stops at ${}^4\text{He}$ due to very small β . However, in the model considered here β is much larger than the canonical value, even being close or exceeding unity. In such conditions much heavier primordial elements can be produced [22]. It is possible that stars which initiated with more metals than the usual ones could look older than they are and if their age is evaluated by the standard nuclear chronology, they might even look older than the universe.

5 Conclusion

The scenario may be speculative but not too unnatural and explains a lot:

1. Superheavy BH and early quasar formation with plenty of metals around.
2. High abundance of supernovae and gamma-bursts at $z \gg 1$.
3. Existence very old stars in the Galaxy and very old galaxies.

Additionally, new types of stellar-like objects from the very early universe and probably abundant cosmic antimatter in the Galaxy are predicted [23]. A study of astrophysics of such new kind of stars is in order.

Acknowledgement. This work was supported by the grant of the Russian Federation government 11.G34.31.0047.

References

- [1] A.D. Dolgov, S.I. Blinnikov, Phys.Rev. D89 (2014) 021301; arXiv:1409.5736 [astro-ph.HE].
- [2] A. Dolgov, V.Halenska, I.Tkachev, JCAP 1410 (2014) 10, 047; arXiv:1406.2445 [astro-ph.CO].

- [3] Planck Collaboration, arXiv:1303.5076 [astro-ph.CO].
- [4] J.J. Cowan, C. Sneden, S. Burles, *et al* Ap.J. 572 (2002) 861, astro-ph/0202429.
- [5] J. Kalirai, Nature 486 (2012) 90, arXiv:1205.6802.
- [6] A. Frebe, N. Christlieb, J.E. Norris, C. Thom Astrophys. J. 660 (2007) L117; astro-ph/0703414.
- [7] H. E. Bond, E. P. Nelan, D. A. VandenBerg, G. H. Schaefer, D. Harmer, Astrophys. J. Lett. 765, L12 (2013); arXiv:1302.3180.
- [8] W. Zheng, M. Postman, A. Zitrin, *et al*, arXiv:1204.2305.
- [9] D. Coe, A. Zitrin, M. Carrasco, *et al*, Astrophys. J. 762 (2013) 32; e-Print: arXiv:1211.3663.
- [10] F. Melia, arXiv:1403.0908 [astro-ph.CO].
- [11] Daniel J. Mortlock, *et al*, Nature 474 (2011) 616, arXiv:1106.6088.
- [12] R. C. E., Gebhardt, K., Gültekin, K., *et al*. 2012, Nature, 491, 729.
- [13] A. Seth, R van den Bosch, S. Mieske, *et al*, arXiv:1409.4769 [astro-ph.GA].
- [14] A. Cucchiara, A.J. Levan, D.B. Fox, N.R. Tanvir, T.N. Ukwatta, E. Berger, T. Kruhler, A.Kupcu Yoldas, X.F. Wu, *et al* Astrophys. J. 736 (2011) 7; e-Print: arXiv:1105.4915 [astro-ph.CO] .
- [15] J. Strader, A. C. Seth, D. A. Forbes, *et al*, Astrophys. J. Lett. 775, L6 (2013), arXiv:1307.7707.
- [16] A. Dolgov, J. Silk, Phys. Rev. D 47 (1993) 4244;
A. D. Dolgov, M. Kawasaki, N. Kevlishvili, Nuclear Physics B 807 (2009) 229, arXiv:0806.2986.
- [17] I. Affleck, M. Dine, Nuclear Physics B 249 (1985) 361
- [18] S.R. Coleman, E.J. Weinberg, Phys.Rev. D7 (1973) 1888.
- [19] F. Ozel, D. Psaltis, R. Narayan, J.E. McClintock, arXiv:1006.2834v2 [astro-ph.GA],
- [20] L. Kreidberg, C.D. Bailyn, W.M. Farr, V. Kalogera, arXiv:1205.1805v1 [astro-ph.HE].
- [21] W.M. Farr, N. Sravan, A. Cantrell, L. Kreidberg, C.D. Bailyn, I. Mandel, V. Kalogera, arXiv:1011.1459v2 [astro-ph.GA]
- [22] R. Nakamura, M.-A. Hashimoto, S.-I. Fujimoto, N. Nishimura, K. Sato, arXiv:1007.0466;
S. Matsuura, S.-I. Fujimoto, S. Nishimura, M.-A. Hashimoto, K. Sato, Phys. Rev. D 72, 123505 (2005), arXiv:astro-ph/0507439;
S. Matsuura, S.-I. Fujimoto, M.-A.Hashimoto, K. Sato, Phys. Rev. D 75, 068302 (2007), arXiv:0704.0635;
S. Matsuura, A. D. Dolgov, S. Nagataki, K. Sato, Progress of Theoretical Physics 112 (2004) 971, arXiv:astro-ph/0405459.
- [23] C. Bambi and A. D. Dolgov, Nuclear Physics B, 784 (2007) 132, arXiv:astro-ph/0702350;
S.I. Blinnikov, A.D. Dolgov, K.A. Postnov, arXiv:1409.5736 [astro-ph.HE].

From the references, the author appears to be unaware of HGD cosmology and its strong constraints on early star formation due to high kinematic viscosities existing in the plasma epoch.

"Within the errors, the age of HD 140283 does not conflict with the age of the Universe, 13.77 +/- 0.06 Gyr, based on the microwave background and Hubble constant, but it must have formed soon after the big bang." Quoting from Bond et al., ref. 7.